

各推薦地の遺存固有種・新固有

分類群		学名	奄美・琉球の固有性
			◎:遺存固有 ○:新固有
哺乳類	トゲネズミ類 アマミトゲネズミ トクノシマトゲネズミ オキナワトゲネズミ	<i>Tokudaia osimensis</i> <i>Tokudaia tokunoshimensis</i> <i>Tokudaia muenninki</i>	◎○
	ケナガネズミ	<i>Diplothrix legata</i>	◎
	アマミノクロウサギ	<i>Pentalagus furnessi</i>	◎
鳥類	ルリカケス	<i>Garrulus lidthi</i>	◎

分類群		学名	奄美・琉球の固有性
			◎:遺存固有 ○:新固有
爬虫類	リュウキュウヤマガメ	<i>Geoemyda japonica</i>	◎
爬虫類	クロイトカゲモドキ類 クロイトカゲモドキ オビトカゲモドキ イヘヤトカゲモドキ マダラトカゲモドキ クメトカゲモドキ	<i>Goniurosaurus kuroiuae kuroiuae</i> <i>Goniurosaurus kuroiuae splendens</i> <i>Goniurosaurus kuroiuae toyamai</i> <i>Goniurosaurus kuroiuae orientalis</i> <i>Goniurosaurus kuroiuae yamashinae</i>	◎○
爬虫類	オキナワヤモリ	<i>Gekko sp.</i>	◎
爬虫類	キノボリトカゲ類 オキナワキノボリトカゲ サキシマキノボリトカゲ ヨナグニキノボリトカゲ	<i>Japalura polygonata polygonata</i> <i>Japalura polygonata ishigakiensis</i> <i>Japalura polygonata donan</i>	○
爬虫類	バーバートカゲ	<i>Plestiodon barbouri</i>	◎
爬虫類	ミヤコカナヘビ	<i>Takydromus toyamai</i>	◎

分類群		学名	奄美・琉球の固有性
			◎:遺存固有 ○:新固有
爬虫類	アオヘビ類 サキシマアオヘビ リュウキュウアオヘビ	<i>Cyclophiops herminae</i> <i>Cyclophiops semicarinatus</i>	○
爬虫類	キクザトサワヘビ	<i>Opisthotropis kikuzatoi</i>	◎
爬虫類	ヒバア類 ガラスヒバア ヤエヤマヒバア ミヤコヒバア	<i>Amphiesma pryeri</i> <i>Amphiesma ishigakiense</i> <i>Amphiesma conelarum</i>	○
爬虫類	ミヤコヒメヘビ	<i>Calamaria pfefferi</i>	◎
爬虫類	アカマタ	<i>Dinodon semicarinatum</i>	◎
爬虫類	ハイ類 ハイ クメジマハイ ヒヤン	<i>Sinomicrurus japonicus boettgeri</i> <i>Sinomicrurus japonicus takarai</i> <i>Sinomicrurus japonicus japonicus</i>	◎○
爬虫類	ヒメハブ	<i>Ovophis okinavensis</i>	◎

分類群		学名	奄美・琉球の固有性
			◎:遺存固有 ○:新固有
両生類	シリケンイモリ	<i>Cynops ensicauda</i>	◎
両生類	イボイモリ	<i>Echinotriton andersoni</i>	◎
両生類	ミヤコヒキガエル	<i>Bufo gargarizans miyakonis</i>	◎
両生類	ハロウエルアマガエル	<i>Hyla hallowellii</i>	◎
両生類	リュウキュウアカガエル アマミアカガエル	<i>Rana ulma</i> <i>Rana kobai</i>	◎○
両生類	ナミエガエル	<i>Limnonectes namiyei</i>	◎
両生類	イシカワガエル類 オキナワイシカワガエル アマミイシカワガエル	<i>Odorrana splendida</i> <i>Odorrana ishikawae</i>	◎○

分類群		学名	奄美・琉球の固有性
			◎:遺存固有 ○:新固有
両生類	ハナサキガエル類 アマミハナサキガエル ハナサキガエル オオハナサキガエル	<i>Odorrana amamiensis</i> <i>Odorrana narina</i> <i>Odorrana supranarina</i>	○
両生類	オットンガエル ホルストガエル	<i>Babina subaspera</i> <i>Babina holsti</i>	◎○
両生類	オキナワアオガエル アマミアオガエル	<i>Rhacophorus viridis viridis</i> <i>Rhacophorus viridis amamiensis</i>	◎○
昆虫類	ヤンバルテナゴコガネ	<i>Cheirotonus jambar</i>	◎
	マルバネクワガタ類 アマミマルバネクワガタ ウケジマルバネクワガタ(亜種) オキナワマルバネクワガタ ヤエヤママルバネクワガタ ヨナグニマルバネクワガタ(亜種)	<i>Neolucanus protogenetivus</i> <i>protogenetivus</i> <i>Neolucanus protogenetivus hamaii</i> <i>Neolucanus okinawanus</i> <i>Neolucanus insulicola insulicola</i> <i>Neolucanus insulicola donan</i>	◎○

分類群		学名	奄美・琉球の固有性
			◎:遺存固有 ○:新固有
甲殻類	サワガニ類		
	アマミクサワガニ属		
	アマミナミサワガニ	Amamiku amamensis	
	カクレサワガニ	Amamiku occulta	
	ミナミサワガニ属		
	オキナワミナミサワガニ	Candidiopotamon okinawense	
	クメジマミナミサワガニ	Candidiopotamon kumejimense	
	トカシキミナミサワガニ	Candidiopotamon tokashikense	
	サワガニ属		
	サカモトサワガニ	Geothelphusa sakamotoana	
	リュウキュウサワガニ	Geothelphusa obtusipes	◎○
	ヒメユリサワガニ	Geothelphusa tenuimanus	
	アラモトサワガニ	Geothelphusa aramotoi	
	ケラマサワガニ	Geothelphusa amagui	
	オキナワオオサワガニ	Geothelphusa grandiovata	
	イヘヤオオサワガニ	Geothelphusa iheya	
	トカシキオオサワガニ	Geothelphusa levicervix	
	クメジマオオサワガニ	Geothelphusa kumejima	
	ムラサキサワガニ	Geothelphusa marginata	
	カシヨクサワガニ	Geothelphusa fulva	
ミネイサワガニ	Geothelphusa minei		
Ryukyum属			
ヤエヤマママガニ	Ryukyum yaeyamense		

## の状態でみられる生物種と文献のリスト

## 文献

## ■遺存固有性

Suzuki, H., Tsuchiya, K., and Takezaki, N. 2000. A molecular phylogenetic framework for the ryukyu endemic rodents *Tokudaia osimensis* and *Diplothrix legata*. *Molecular Phylogenetics and Evolution*. 15 (1) 15-24.

Sato, J.J, and Suzuki, H. 2004. Phylogenetic relationships and divergence times of the genus *Tokudaia* within Murinae (Muridae; Rodentia) inferred from the nucleotide sequences encoding the mitochondrial cytochrome b gene and nuclear recombination-activating gene 1 and interphotoreceptor retinoid-binding protein. *Canadian Journal of Zoology* 82: 1343-1351.

## ■新固有性

Endo, H. and Tsuchiya, K. 2006. A new species of Ryukyu spiny rat, *Tokudaia* (Muridae: Rodentia), from Tokunoshima Island, Kagoshima Prefecture, Japan. *Mammal Study*, 31 (1) : 47-57.

Kaneko, Y. 2000. Morphological discrimination of the Ryukyu spiny rat (genus *Tokudaia*) between the islands of Okinawa and Amami Oshima, in the Ryukyu Islands, southern Japan. *Mammal Study*, 26 (1) :17-33.

Suzuki, H., Masahiro A. Iwasa, M. A., Ishii, N., Nagaoka, H. and Tsuchiya, K. 1999. The genetic status of two insular populations of the endemic spiny rat *Tokudaia osimensis* (Rodentia, Muridae) of the Ryukyu Islands, Japan. *Mammal Study*. 24 (1) : 43-50

土屋公幸・若菜茂晴・鈴木仁・服部正策・林良博. 1989. トゲネズミの分類学的研究 I. 遺伝的分化. 国立科学博物館専報, 22 : 227-234.

Suzuki, H., Tsuchiya, K., and Takezaki, N. 2000. A molecular phylogenetic framework for the ryukyu endemic rodents *Tokudaia osimensis* and *Diplothrix legata*. *Molecular Phylogenetics and Evolution*. 15 (1) 15-24

Sato, J.J, and Suzuki, H. 2004. Phylogenetic relationships and divergence times of the genus *Tokudaia* within Murinae (Muridae; Rodentia) inferred from the nucleotide sequences encoding the mitochondrial cytochrome b gene and nuclear recombination-activating gene 1 and interphotoreceptor retinoid-binding protein. *Canadian Journal of Zoology* 82: 1343-1351.

Yamada, F., Takaki, M. and Suzuki, H. 2002. Molecular phylogeny of Japanese Leporidae, the Amami rabbit *Pentalagus furnessi*, the Japanese hare *Lepus brachyurus*, and the mountain hare *Lepus timidus*, inferred from mitochondrial DNA sequences. *Genes Genet. Syst.* 77:107-116.

Matthee, C.A., B. J. Vuuren, D. Bell, & T. J. Robinson. 2004. A Molecular supermatrix of the rabbits and hares (Leporidae) allows for the identification of five international exchange during the Miocene. *Sytematic Biology*. 53 : 433-447.

山階芳麿. 1942. 琉球列島特産鳥類3種の分類学的位置と生物地理学的意義について. *Trans. Biogeogr. Soc. Japan* 3 : 319-328.

梶田学, 川路則友, 山口恭弘, Khan AA. 1999. ムリカケス *Garrulus lidthi* の系統関係について—DNA解析と形態の両面から—. 日本鳥学会1999年度退会講演要旨集. p44, 東京大学.

Akimova, A. Haring, E., Kryukov, S., Kryukov, A. 2007. First insights into a DNA sequence based phylogeny of the Eurasian Jay *Garrulus glandarius*. *Рус. о р н и т о л. ж у р н.* 2007. Т о м 16. Э к с п р е с с - в ы п у с к No. 356

文献

■系統解析

Gong, S., H. Shi, Y. Mo, M. Auer, M. Vargas-Ramírez, A. K. Hundsdoerfer, and U. Fritz. 2009. Phylogeography of the endangered black-breasted leaf turtle (*Geoemyda spengleri*) and conservation implications for other chelonians. *Amphibia-Reptilia* 30: 57-62.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189-204.

■系統解析

Honda, M., T. Kurita, M. Toda, and H. Ota. 2014. Phylogenetic relationships, genetic divergence, historical biogeography and conservation of an endangered gecko, *Goniurosaurus kuroiwae* (Squamata: Eublepharidae), from the Central Ryukyus, Japan. *Zool. Sci.* 30:309-320.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189-204.

■系統解析

Toda, M., Hikida, T. & Ota, H. 2001. Discovery of sympatric cryptic species within *Gekko hokouensis* (Gekkonidae: Squamata) from the Okinawa Islands, Japan, by use of allozyme data. *Zoologica scripta* 30 : 1-11.

■系統解析

Ota, H. 1991. Taxonomic redefinition of *Japalura swinhonis* Günther (Agamidae: Squamata), with a description of a new subspecies of *J. polygonata* from Taiwan. *Herpetologica* 47: 280-294.

Ota, H. 2003. A new subspecies of the Agamid Lizard, *Japalura polygonata* (Hallowell, 1861) (Reptilia : Squamata), from Yonagunijima Island of the Yaeyama Group, Ryukyu Archipelago. *Cur. Herpetol.* 22: 61-71.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189-204.

■系統解析

Brandley, M.C., Ota, H., Hikida, T., Nieto-Montes de Oca, A., Fera-Ortiz, M., Guo, X., Wang, Y. 2012. The phylogenetic systematics of blue-tailed skinks (Plestiodon) and the family Scincidae. *Zoological Journal of the Linnean Society*. 165(1): 163-189.

Honda, M., T. Okamoto, T. Hikida and H. Ota, 2008. Molecular phylogeography of the endemic five-lined skink (*Plestiodon marginatus*) (Reptilia: Scincidae) of the Ryukyu Archipelago, Japan, with special reference to the relationship of a northern Tokara population. *Pacific Sci.*, 62: 351-362.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189-204.

■系統解析

Ota, H., M. Honda, S. Chen, T. Hikida, S. Panha, H. Oh, and M. Matsui. 2002. Phylogenetic relationships, taxonomy, character evolution and biogeography of the lacertid lizards of the genus *Takydromus* (Reptilia : Squamata): a molecular perspective. *Biological Journal of the Linnean Society*. 76: 493-509.

Lin, SM., CA Chen, KY Lue. 2002. Molecular phylogeny and biogeography of the grass lizards genus *Takydromus* (Reptilia: Lacertidae) of East Asia. *Molecular Phylogenetics and Evolution* 22:276-288.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189-204.



文献

■系統解析

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Ota, H. and S. Iwanaga, 1997. A systematic review of the snakes allied to *Amphiesma pryri* (Boulenger) (Squamata: Colubridae) in the Ryukyu Archipelago, Japan. *Zoological Journal of the Linnean Society*, 121: 339–360. **未入手**

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Slowinski JB, Boundy J, Lawson R. 2001. The phylogenetic relationships of Asian coral snakes (Elapidae: Calliophis and Maticora) based on morphological and molecular evidence. *Herpetologica* 57: 233–245.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Malhotra A, Thorpe RS. 2004a. A phylogeny of four mitochondrial gene regions suggests a revised taxonomy for Asian pit vipers (*Trimeresurus* and *Ovophis*). *Molecular Phylogenetics and Evolution* 32 (1): 83–100.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

文献

■系統解析

Tominaga, A., H. Ota, M. Matsui. 2010. Phylogeny and phylogeography of the sword-tailed newt, *Cynops ensicauda* (Amphibia: Caudata), as revealed by nucleotide sequences of mitochondrial DNA. *Molecular Phylogenetics and Evolution* 54:3, 910–921.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Honda M, Matsui M, Tominaga A, Ota H, Tanaka S. 2012. Phylogeny and biogeography of the Anderson's crocodile newt, *Echinotriton andersoni* (Amphibia: Caudata), as revealed by mitochondrial DNA sequences. *Mol Phylogenet Evol* 65: 642–653.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Riehl CJ, Lell JT, Maxson LR. 1995. Relationships among Palearctic Hyla: insights from immunology. *Biochemical Systematics and Ecology* 23: 245–249. **未入手**

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Matsui M. 2011. On the brown frogs from the Ryukyu Archipelago, Japan, with descriptions of two new species (Amphibia, Anura). *Curr Herpetol* 30: 111–128.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Emerson, S. B., and D. Berrigan, 1993. Systematics of Southeast Asian ranids : multiple origins of voicelessness in the subgenus *Limnonectes* (Fitzinger). *Herpetologica*, 49 : 22–31. **未入手**

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Kuramoto, M., Satou, N., Oumi, S., Kurabayashi, A. & Sumida, M. 2011. Inter- and intra-island divergence in *Odorrana ishikawae* (Anura, Ranidae) of the Ryukyu Archipelago of Japan, with description of a new species. *Zootaxa*, 2767, 25–40.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

文献

■系統解析

Matsui M, Shimada T, Ota H, Tanaka-Ueno T (2005b) Multiple invasions of the Ryukyu Archipelago by Oriental frogs of the subgenus *Odorrana* with phylogenetic reassessment of the related subgenera of the genus *Rana*. *Mol Phylogenet Evol* 37: 733–742.

Matsui, M. 1994. A taxonomic study of the *Rana narina* complex, with description of three new species (Amphibia: Ranidae). *Zool. J. Linn. Soc.* 111:385–415.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

Tominaga, A., M. Matsui, and K. Nakata. 2014. Genetic diversity and differentiation of a Ryukyu endemic frog *Babina holsti* as revealed by mitochondrial DNA. *Zool. Sci.* 31: 64–70.

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

■系統解析

■総説

Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*. 40: 189–204.

細谷忠嗣, 荒谷邦雄. 論文執筆中。

細谷忠嗣, 荒谷邦雄. 2010. DNAによる系統解析の紹介. *コガネ博2010レジュメ集*. pp.15–18.

細谷忠嗣, 荒谷邦雄, 七里浩志, 佐々木健志, 当山昌直. 2006. ミトコンドリア遺伝子からみたテナガコガネ亜科の系統関係とヤンバルテナガコガネの遺伝的多様性について. *日本鞘翅学会第19回大会講演要旨集*.

細谷忠嗣, 荒谷邦雄, 澤志泰正, 佐々木健志, 当山昌直. 2005. ミトコンドリア 16S rRNA 遺伝子からみたテナガコガネ亜科の系統関係. *日本鞘翅学会第18回大会講演要旨集*.

細谷忠嗣, 荒谷邦雄. 2006. 琉球列島におけるマルバネクワガタ属の分子生物地理. *昆虫と自然*. 41(4) : 5–10.

文献

瀬川涼子. 2011. サワガニ類の分子系統学的研究. 川井唯史・中田和義(編). エビ・カニ・ザリガニ—淡水甲殻類の保全と生物学. 生物研究社. pp103-110.

瀬川涼子. 2000. 琉球列島のサワガニ属の分子系統学的研究. 月刊海洋総特集 分子海洋学—分子生態学と海洋学の接点. 32(4): 241-245.

諸喜田茂光. 1996. 琉球列島の河川陸封コエビ類とサワガニ類の起源論. 地学雑誌. 105(3) 343-353.